

***Mirohelminthosporium* gen. nov. for an atypical *Helminthosporium* species and *H. matsushimae* nom. nov.**

KAI ZHANG¹, HE ZHANG², DE-WEI LI^{3,4}, RAFAEL F. CASTAÑEDA-RUIZ^{5*}

¹ Shandong Agriculture and Engineering University,
Jinan, Shandong 250100, China

² Department of Landscaping, Shandong Yingcai University,
Jinan, Shandong 250104, China

³ The Connecticut Agricultural Experiment Station, Valley Laboratory,
153 Cook Hill Road, Windsor, CT 06095, USA

⁴ Co-Innovation Center for Sustainable Forestry in Southern China,
Nanjing Forestry University, Nanjing, Jiangsu 210037, China

⁵ Instituto de Investigaciones Fundamentales en Agricultura Tropical Alejandro de Humboldt
(INIFAT), Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

* CORRESPONDENCE TO: rffcastanedaruiz@gmail.com

ABSTRACT—*Mirohelminthosporium* is proposed as a new genus for *Helminthosporium bigenum*, characterized by polytretic and blastic conidial ontogeny on the apical conidiogenous cells. A replacement name, *Helminthosporium matsushimae*, is proposed for the illegitimate later homonym *H. cylindrosporum* Matsush.

KEY WORDS—asexual Ascomycota, *Helminthosporium dimorphosporum*, hyphomycetes, systematics

Introduction

Helminthosporium Link typified by *H. velutinum* Link is characterized by macronematous unbranched erect conidiophores (Link 1809). The conidiogenous cells are polytretic, integrated, terminal and intercalary, and determinate, with the conidiogenous loci at the apex and laterally below the septa. The conidia are solitary, obclavate to cylindrical-obclavate, distoseptate,

sometimes rostrate, smooth, subhyaline, olivaceous, brown, and usually with an evident dark brown or brown basal scar. The *Helminthosporium* generic concept treated by Alcorn (1983, 1988), Ellis (1961, 1971), Hughes (1958, 1980a,b), Shirouzu & Harada (2004), Voglmayr & Jaklitsch (2017), and Zhang & Zhang (2009) clarified the morphological and molecular phylogenetic delimitation of the genus. Additionally, several *Helminthosporium* species with blastocatenate distoseptate and dictyoseptate conidia were transferred into *Ellismasporium* R.F. Castañeda & X.G. Zhang and *Stanhughesiella* R.F. Castañeda & D.W. Li (Castañeda-Ruiz & al. 2017).

Helminthosporium bigenum has an unusual combination of holoblastic euseptate conidia from apical conidiogenous cells and tretic (enteroblastic) distoseptate conidia from intercalary conidiogenous cells (Matsushima 1993). Conidial ontogeny and conidiogenous events have been considered the fundamental criteria for asexual fungal taxonomy and generic delimitation (Kendrick 2017, Seifert & al. 2011). Consequently, we propose the new genus *Mirohelminthosporium* to accommodate the atypical *Helminthosporium* species, *H. bigenum*. A replacement name, *Helminthosporium matsushimae*, is proposed for the illegitimate later homonym *H. cylindrosporum* Matsush.

Taxonomy

Mirohelminthosporium K. Zhang, D.W. Li & R.F. Castañeda, **gen. nov.**

MB 833028


Differs from *Helminthosporium* by its holoblastic and primarily monoblastic conidial ontogeny, and its production of euseptate conidia on the apical conidiogenous cells beside or near the tretic loci.

TYPE SPECIES: *Helminthosporium bigenum* Matsush. [= *Mirohelminthosporium bigenum* (Matsush.) K. Zhang & al.]

ETYMOLOGY: Latin, *Mirus*- meaning wonderful, extraordinary, remarkable + Latin, *-helminthosporium*, referring to the genus *Helminthosporium*.

COLONIES on CMA medium hairy, brown to dark brown. CONIDIOPHORES macronematous, mononematous, unbranched, erect, straight, cylindrical, septate, smooth, brown to dark brown. CONIDIOGENOUS CELLS biontogenous, sometimes on the same cell: i) polytretic, integrated, mostly intercalary, cylindrical, sometimes terminal with subapical tretic loci and apical blastic loci; ii) monoblastic, integrated, terminal, sometimes with

FIG. 1. *Mirohelminthosporium bigenum* (Adapted from Matsushima 1993, figs 693–696; *H. bigenum* holotype, MFC OP-440). A. Conidia from blastic loci. B. Conidia from tretic loci. C. Conidiogenous cells.



several tretic loci near below the blastic locus on the same cell. CONIDIA solitary, dimorphic: i) acropleurogenous, mostly pleurogenous, cylindrical, oblong, straight or curved, rounded at the ends, melanized or obscure at the base, distoseptate, smooth, brown to dark brown or black, dry, borne on tretic conidiogenous loci; ii) acrogenous, solitary, cylindrical, clavate to subclavate, attenuate, rounded or obtuse at the apex, truncate at the base, euseptate, smooth, brown to dark brown or black, dry, borne on the apical blastic conidiogenous loci.

Mirohelminthosporium bigenum (Matsush.) K. Zhang, D.W. Li &

R.F. Castañeda, **comb. nov.**

FIG. 1

MB 833029

= *Helminthosporium bigenum* Matsush., Matsush. Mycol. Mem.7: 53 (1993).

Helminthosporium matsushimae D.W. Li, K. Zhang &

R.F. Castañeda, **nom. nov.**

FIG. 2

MB 833030

= *Helminthosporium cylindrosporum* Matsush., Matsush. Mycol.

Mem.7: 54 (1993), nom. illeg. (non Sacc. 1876).

NOTES: Index Fungorum (2019) lists 770 taxonomic names in *Helminthosporium*, but many of these have been transferred to the genera *Bipolaris* Shoemaker, *Curvularia* Boedijn, *Drechslera* S. Ito, *Exserohilum* K.J. Leonard & Suggs, *Exosporium* Link, *Ellismarsporium*, and several other genera.

Anisospadicoides R.F. Castañeda & al. is another genus with blastic and tretic conidial ontogeny but in that instance, both types of conidiogenesis produce euseptate conidia (Qiao & al. 2019).

Helminthosporium dimorphosporum Hol.-Jech. (FIG. 3), which produces distoseptate and euseptate conidia through the same tretic loci (Holubová-Jechová 1987), does not fit the *Helminthosporium* generic concept and needs to be revised.

Acknowledgments

This work was financed by the National Natural Science Foundation Program of PR China (31870016). The authors express their sincere gratitude to Dr. Josiane S. Monteiro and Dr. Flavia Rodrigues Barbosa for their critical review of the manuscript. We acknowledge the facilities provided by Dr. P.M. Kirk and Dr. K. Bensc through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.




FIG. 2. *Helminthosporium matsushimae*
(Adapted from Matsushima 1993, figs 239, 686; *H. cylindrosporum* holotype, MFC OP-658).
A. Conidia. B. Conidiogenous cells.




FIG. 3. *Helminthosporium dimorphosporum*
(Adapted from Holubová-Jechová 1987, fig. 2:3, *H. dimorphosporum* holotype, PRM 842 738).
A. Distoseptate conidia. B. Euseptate conidia. C. Conidiogenous cells.

Literature cited

- Alcorn JL. 1983. Generic concepts in *Drechslera*, *Bipolaris* and *Exserohilum*. Mycotaxon 17: 1–86.
- Alcorn JL. 1988. The taxonomy of ‘*Helminthosporium*’ species. Annual Review of Phytopathology 26: 37–56. <https://doi.org/10.1146/annurev.py.26.090188.000345>
- Castañeda-Ruiz RF, Li DW, Zhang XG, Kendrick B, Ramos-García B, Pérez-Martínez S, Sosa D. 2017. *Ellismarsporium* gen. nov. and *Stanhughesiella* gen. nov. to accommodate atypical *Helminthosporium* and *Corynesporella* species. Mycotaxon 132(4): 759–766. <https://doi.org/10.5248/132.759>
- Ellis, M.B. 1961. Dematiaceous hyphomycetes. III. Mycological Paper 82. 55 p.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England. 608 p.
- Holubová-Jechová V. 1987. Studies on hyphomycetes from Cuba VI. New and rare species with tetric and phialidic conidiogenous cells. Česká Mykologie 41: 107–114.
- Hughes SJ. 1958. Revisiones hyphomycetum aliquot cum appendice de nominibus rejiciendis. Canadian Journal of Botany 36(6): 727–836. <https://doi.org/10.1139/b58-067>
- Hughes SJ. 1980a. *Helminthosporium velutinum*. Fungi Canadenses 163: 1–2.
- Hughes SJ. 1980b. New Zealand Fungi 27. New species of *Guedea*, *Hadrosporium*, and *Helminthosporium*. New Zealand Journal of Botany 18: 65–72. <https://doi.org/10.1080/0028825X.1980.10427233>
- Index Fungorum. 2019. www.indexfungorum.org./Names/Names.asp (accessed 20 Sept. 2019).
- Kendrick B. 2017. The fifth kingdom (4th ed). Hackett publishing company. 416 p.
- Link HF. 1809. Observationes in ordines plantarum naturales. Dissertatio I. Magazin der Gesellschaft Naturforschenden Freunde Berlin 3(1): 3–42.
- Matsushima T. 1993. Matsushima Mycological Memoirs no. 7. Matsushima Fungus Collection, Kobe, Japan.
- Qiao M, Li DW, Yu ZF, Zhang K, Castañeda-Ruiz RF. 2019. *Spadicoides matsushimae* sp. nov., and *Anisospadicoides* gen. nov. for two atypical *Spadicoides* species. Mycotaxon 134: 161–167. <https://doi.org/10.5248/134.161>
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.
- Shirouzu T, Harada Y. 2004. Notes on species of *Helminthosporium* and its allied genera in Japan. Mycoscience 45: 17–23. <https://doi.org/10.1007/S10267-003-0146-8>
- Voglmayr H, Jaklitsch WM. 2017. *Corynespora*, *Exosporium* and *Helminthosporium* revisited – new species and generic reclassification. Studies in Mycology 87: 43–76. <https://doi.org/10.1016/j.simyco.2017.05.001>
- Zhang M, Zhang TY. 2009. Taxonomic studies of *Helminthosporium* from China 4. Six new species and a key to *Helminthosporium* from China. Mycotaxon 109: 399–413. <https://doi.org/10.5248/109.399>